

Quantum Groups, Strings and HTSC materials

Sher Alam¹, M. O. Rahman², T. Yanagisawa¹, and H. Oyanagi¹

¹*Physical Science Division, ETL, Tsukuba, Ibaraki 305, Japan*

²*GUAS & Photon Factory, KEK, Tsukuba, Ibaraki 305, Japan*

Abstract

Previously we have indicated the relationship between quantum groups and strings via WZWN models. In this note we discuss this relationship further and point out its possible applications to cuprates and related materials. The connection between quantum groups and strings is one way of seeing the validity of our previous conjecture [i.e. that a theory for cuprates may be constructed on the basis of quantum groups]. The cuprates seems to exhibit statistics, dimensionality and phase transitions in novel ways. The nature of excitations [i.e. quasiparticle or collective] must be understood. The Hubbard model captures some of the behaviour of the phase transitions in these materials. On the other hand the phases such as stripes in these materials bear relationship to quantum group or string-like solutions. One thus expects that the relevant solutions of Hubbard model may thus be written in terms of stringy solutions. In short this approach may lead to the non-perturbative formulation of Hubbard and other condensed matter Hamiltonians. The question arises that how a 1-d based symmetry such as quantum groups can be relevant in describing a 3-d [spatial dimensions] system such as cuprates. The answer lies in the key observation that strings which are 1-d objects can be used to describe physics in d dimensions. For example gravity [which is a 3-d [spatial] plus time] phenomenon can be understood in terms of 1-d strings. Thus we expect that 1-d quantum group object induces physics in 2-d and 3-d which may be relevant to the cuprates.

In a previous work one of us [1] has advanced the conjecture that one should attempt to model the phenomena of antiferromagnetism and superconductivity by using quantum symmetry group. Following this conjecture to model the phenomena of antiferromagnetism and superconductivity by quantum symmetry groups, three toy models were proposed [2], namely, one based on $SO_q(3)$ the other two constructed with the $SO_q(4)$ and $SO_q(5)$ quantum groups. Possible motivations and rationale for these choices were outlined [2]. In [3] a model to describe quantum liquids in transition from 1d to 2d dimensional crossover using quantum groups was outlined. In [5] the classical group $SO(7)$ was proposed as a toy model to understand the connections between the competing phases and the phenomenon of pseudo-gap in High Temperature Superconducting Materials [HTSC]. Then we proposed in [6] an idea to construct a theory based on patching critical points so as to simulate the behavior of systems such as cuprates. To illustrate our idea we considered an example discussed by Frahm et al., [4]. The model deals with antiferromagnetic spin-1 chain doped with spin-1/2 carriers. In [7] the connection between Quantum Groups and 1-dimensional [1-d] structures such as stripes was outlined. The main point of [7] is to emphasize that *1-d structures play an important role in determining the physical behaviour [such as the phases and types of phases these materials are capable of exhibiting] of cuprates and related materials.*

The question arises that how a 1-d based symmetry such as quantum groups can be relevant in describing a 3-d [spatial dimensions] system such as cuprates. *The answer lies in the key observation that strings which are 1-d objects can be used to describe physics in d dimensions.* For example gravity [9] [which is a 3-d [spatial] plus time] phenomenon can be understood in terms of 1-d strings. Thus we expect that 1-d quantum group object induces physics in 2-d and 3-d which may be relevant to the cuprates. However we like to point out that the notion of string is more general and does not have to be restricted to quantum groups. The reason for our choice are several. For one quantum groups are intimately

connected with WZWN models and to strings. In turn WZWN based models are relevant to disordered systems. The exact solution of Hubbard model in 1-d is governed by quantum group symmetry. In turn the Hubbard model captures much of the physics of cuprates. We note that the t-J model and its various generalizations are also advocated for the study of cuprates. However the t-J model is but a limit of Hubbard model [i.e. when $U/t \rightarrow \infty$]. Quantum critical points may be naturally understood in terms of quantum groups.

Let us briefly comment on three different general aspects of the cuprates and related materials:

- Statistics and fractionalization:-The cuprates seem to exhibit exotic statistics and charge fractionalization. Indications for electron fractionalization from Angle-Resolved Photoemission Spectroscopy [ARPES] have been reported. We have suggested to measure fractionalization by using SET based experiments [8] *. We note that the Fermi liquid is characterized by sharp fermionic quasiparticle excitations and has a discontinuity in the electron momentum distribution function. In contrast the Luttinger liquid is characterized by charge e spin 0 bosons and spin 1/2 charge 0 and the fermion is a composite of these [i.e. fractionalization]. It is well-known that transport properties

*After about a week or so later of our submission a paper by Sentil and Fisher “Detecting fractions of electrons in the high- T_c cuprates” cond-mat/0011345, appeared without reference to our work. Some mathematical details were worked out but it used or was in line with basically our idea. What follows is the opinion of Sher Alam: I [Sher Alam] informed by email Sentil of this situation, in his arrogance he never replied. Such an attitude highly annoys me. We all use each other ideas in some form or the other, but more modest people have the courage to admit it. I hate pretentious and arrogant attitudes. Knowledge is God given right and not the exclusive right of some as they would have us believe it.

are defined via correlation functions. The correlation functions of a Luttinger liquid have a power law decays with exponents that depend on the interaction parameters. Consequently the transport properties of a Luttinger liquid are very different from that of a Fermi liquid. Photoemission experiments on Mott insulating oxides seems to indicate the spinon and holon excitations of a charge Luttinger liquid. However the experimental signatures of Luttinger liquid are not totally convincing. To this end we propose SET based experiments to determine the Luttinger liquid behaviour of the cuprates.

- Mixed dimensionality:- The cuprates and related materials seem to exhibit mixed dimensionality. This can be seen in many materials. We give an example of $\text{Ca}_{2+x}\text{Y}_{2-x}\text{Cu}_5\text{O}_{10}$ [10], a material which simple [i.e. simplicity in structural aspect and controlled hole doping] and exhibits mixed dimensionality and can provide insight into the spin and charge dynamics of more complicated HTSC material. As a consequence of mixed dimensionality this material exhibits ferromagnetism and antiferromagnetism, introduction of holes leads to novel spin-charge dynamics in this magnetically frustrated system.
- Phases:-A variety of phases have been reported in cuprates, for example, antiferromagnetic insulator, superconducting, metallic, strange metal, spin-glass, and insulating.

It has been known that the symmetry of Hubbard model is $SU(2) \times SU(2)/Z_2 \equiv SO(4)$, this can be taken as a motivation to consider a gauge model or NLSM for Hubbard based on $SU(2)$, $SU(2) \times$, $SO(4)$ or even Z_2 or many combinations and generalizations thereof depending on one's point of view. We take the following point of view. It is known that quantum groups in the context of usual HH are tied to 1-d. This is used as an argument against the application of quantum group to higher dimensions. We take the opposite atti-

tude. Using the string notion instead of going from 1-d, to 2-d via the usual route, we assume that fundamental entities are 1-d objects [subject to quantum group symmetry] i.e. strings, for example we introduce the notion of Hubbard string. And attach the known symmetry of Hubbard Hamiltonian i.e. $SU(2) \times SU(2)/Z_2 \equiv SO(4)$ for example as a Chan-Paton factor[†]. We conjecture that the solution of 2-d and 3-d Hubbard Hamiltonian can be written in terms of the 1-d Hubbard string solutions. This conjecture can be tested by looking at specific examples. We thus have provided a general framework for strongly correlated electron systems, such as Quantum Hall systems and HTSC materials and related materials. The recent data supports our earlier conjecture of important role of 1-d systems tied to quantum groups.

P.W.Anderson and others have tried to prove spin charge separation in 2-d. In our framework this is not necessary and follows naturally as a consequence of the Hubbard string. It is clear that spin charge separation takes place on the 1-d Hubbard string, the dynamics of the string leads to spin charge separation in 2-d and restricted confinement as appears in the form of stripes observed in experiment. The ARPES data shows a significant Fermi surface with hotspots, this cannot be easily explained in terms of Fermi Liquid or arranged non Fermi Liquid behaviour. The collective dynamics of 1-d strings can provide a suitable explanation. For example one can interpret that the holes Fermi surface are the regions where the collective or luttinger liquid-like behaviour dominates. Let us try to see this

[†]Chan-Paton factor:-An open string has boundaries, i.e. endpoints, in quantum with distinguished endpoints it is natural to associate degrees of freedom with these in addition to fields propagating in bulk. Moreover it is natural and necessary to have simply boundary conditions. For example in string theory of strong interactions, one introduces $SU(3)$ flavor quantum numbers, the endpoints have quark-antiquark attached connected by ‘color-electric flux tube

point intuitively, string theory has several conservation laws, which follows due to various symmetries. Now some of these symmetries are broken in the target space, for example in context of gravity conformal invariance is broken in the target space by gravity and preserved on the 2-d surface generated by string and yet it is the condition of conformal invariance on string generated surface that gives Einstein's equation coupled to a scalar field in the target space. Thus in an analogous manner one may have obtain a situation where Luttinger liquid symmetry is preserved in some regions of the Fermi surface and badly broken in others. As the Hubbard strings fluctuate they generate a Fermi surface where Luttinger correlations are preserved in only some regions. The area or volume where non-Fermi liquid persists is expected to be proportional to the number of strings, which is turn is determined by physical parameters such as doping etc. A challenging problem is to define precisely a measure for the “correlations” in the system.

In short we expect that by formulating the underling theory in terms of collective excitations such as strings [which are indicative of Luttinger liquids] we can represent a general theory for strongly correlated systems in 2-d and 3-d. Yet another support for our point of view comes from data of Uemura, where he finds 2-d bose like behaviour, and these fit reasonably with Berezinskii-Kosterlitz-Thouless [BKT] type of phase transitions. Within our framework this is natural since conformal invariance is intimately tied with strings. It would be interesting to look into the details of the relationship between Uemura's data and the BKT-type phase transitions.

We now comment briefly on the choice of Hamiltonian and naively on one kind of duality in this context. As already mentioned and as is well-known the t-J model is a special limit of Hubbard model. Keeping this point in mind we want to seek a non-trivial unification of Hubbard Hamiltonian and the t-J model using ‘duality’ of couplings in a new way. To achieve this unification we start with a small step. Let us explain. To this end let us first

recall the usual argument based on perturbation theory which allows us to see the t-J model as the special limit of the Hubbard Hamiltonian [HH]. The HH reads

$$H = -t \sum_{\langle ij \rangle \sigma} c_{i\sigma}^\dagger c_{j\sigma} + U \sum_i n_{i\uparrow} n_{i\downarrow}, \quad (1)$$

where $\langle ij \rangle$ means sum over nearest neighbours, i.e. hybridization between neighbouring atoms. $c_{i\sigma}^\dagger$ ($c_{i\sigma}$) are the creation (annihilation) operators of electrons at site i of spin σ ($\sigma \equiv \uparrow, \downarrow$). An important limit of HH is half-filled and strong coupling, i.e. $U \gg t$.

The intermediate state has one doubly occupied atom and the effective interaction [second-order] is simply

$$H = t \sum_{\langle ij \rangle \sigma'} c_{i\sigma'}^\dagger c_{j\sigma'} \frac{-1}{U} t \sum_{\sigma} c_{i\sigma}^\dagger c_{j\sigma}, \quad (2)$$

which is the Hamiltonian of antiferromagnetically coupled spin- $\frac{1}{2}$ Heisenberg model with coupling $J = 2t^2/U$ per bond. Naively one can regard in some sense the Hamiltonian in Eq. 2 as a ‘dual’ with respect to the coupling U of the Hamiltonian in Eq. 1. This argument is made in the context of degenerate perturbation theory and the assumptions stated above. We now abandon any assumption of perturbation theory and simply assume to start with a Hamiltonian which is the sum of the above two Hamiltonians, i.e.

$$\begin{aligned} H &= -t \sum_{\langle ij \rangle \sigma} c_{i\sigma}^\dagger c_{j\sigma} + U \sum_i n_{i\uparrow} n_{i\downarrow} \\ &\quad + J \sum_{\langle ij \rangle \sigma'} c_{i\sigma'}^\dagger c_{j\sigma'} \sum_{\sigma} c_{i\sigma}^\dagger c_{j\sigma}, \\ J &= -g^2/U, \end{aligned} \quad (3)$$

where g is some suitable coupling.

In conclusion we have indicated new ways of looking at the physics of HTSC and related materials and in general strongly correlated systems. Although in its current form quantum groups are restricted to 1-d, we don’t see this as a problem but a solution. This 1-d

restriction led us to think that we can solve a 2-d or 3-d problem by using string theory as is done in case of gravity. Gravity is a 4-d phenomenon, but can be understood elegantly in terms of string theory [1-d objects]. This approach is different when one tries to go from 1-d to higher dimensions directly. Recent developments in string and topological field theories can further help us to understand the physics of strongly correlated systems.

ACKNOWLEDGMENTS

The Sher Alam's work is supported by the Japan Society for for Technology [JST].

REFERENCES

- [1] Sher Alam, *A Conjecture for possible theory for the description of high temperature superconductivity and antiferromagnetism*. Proceedings of Quantum Phenomena in Advanced Materials at High Magnetic Fields, 4th International Symposium on Advanced Physical Fields [APF-4]. KEK-TH-607, KEK Preprint 98-xxx, cond-mat/9812060.
- [2] Sher Alam, *Quantum Group based Modelling for the description of high temperature superconductivity and antiferromagnetism*, Phys. Lett. A272, (2000), 107-112. KEK-TH-613, KEK Preprint 98-xxx, cond-mat/9903038.
- [3] Sher Alam et al., *Theoretical modeling for quantum liquids from 1d to 2d dimensional crossover using quantum groups*. KEK-TH-619. KEK Preprint 99-xxx, cond-mat/990345.
- [4] Holger. Frahm et al., Phys. Rev. Lett. **81**, (1998), 2116.
- [5] Sher Alam et al., *The Choice of symmetry group for cuprates*, cond-mat/0004269.
- [6] Sher Alam et al., *The patching of critical points using quantum groups*, cond-mat/0004350.
- [7] Sher Alam, *Quantum group conjecture and Stripes* cond-mat/0005168.
- [8] Sher Alam et al., *SET based experiments for HTSC materials* cond-mat/0011037.
- [9] M. Kaku, *Strings, Conformal Fields, and Topology: An Introduction*, Springer-Verlag, 1991.
- [10] H. Yamaguchi et al., Physica. C320, (1999), 167-172.